

NAME: Solutions

EID:

M 427K Exam 2 Version A November 13, 2012 Instructor: James Pascaleff

Problem	Possible	Actual
1	10	
2	7	
3	20	
4	10	
5	5	
6	15	
7	18	
8	15	
Total	100	

INSTRUCTIONS:

- Do all work on these sheets.
- Show all work.
- No books, notes, calculators, or other electronic devices.

HIGHER ORDER EQUATIONS

1. (10 points) Find the general solution of the fourth order equation

$$y^{(4)} - 5y'' + 4y = 0$$

$$0 = r^4 - 5r^2 + 4 = (r^2 - 4)(r^2 - 1) = (r - 2)(r + 2)(r - 1)(r + 1)$$

$$\text{roots: } r_1 = 2, r_2 = -2, r_3 = 1, r_4 = -1$$

$$y = c_1 e^{2t} + c_2 e^{-2t} + c_3 e^t + c_4 e^{-t}$$

POWER SERIES

2. (7 points) Find the radius of convergence of the series

$$L = \lim_{n \rightarrow \infty} \frac{\left| \frac{3^{n+1} x^{n+1}}{(n+1)!} \right|}{\left| \frac{3^n x^n}{n!} \right|} = \lim_{n \rightarrow \infty} \frac{n!}{(n+1)!} \frac{3^{n+1}}{3^n} \frac{|x|^{n+1}}{|x|^n}$$

$$= \lim_{n \rightarrow \infty} \frac{1}{n+1} 3|x| = 0 < 1 \text{ for any } x.$$

So the Radius of convergence is ∞ ; the series converges for all x .

3. (20 points) Seek a power series solution of the following equation at the point $x_0 = 0$:

$$y'' + xy' + 2y = 0$$

Find the recurrence relation, and determine the terms, up to the x^3 term, of the solution that begins with $a_0 = 0$, $a_1 = 1$.

$$y = \sum_{n=0}^{\infty} a_n x^n \quad y' = \sum_{n=0}^{\infty} n a_n x^{n-1}$$

$$xy' = x \sum_{n=0}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} n a_n x^n$$

$$y'' = \sum_{n=0}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$0 = y'' + xy' + 2y = \sum_{n=0}^{\infty} \left[(n+2)(n+1) a_{n+2} + n a_n + 2 a_n \right] x^n$$

$$\Rightarrow 0 = (n+2)(n+1) a_{n+2} + n a_n + 2 a_n = (n+2)(n+1) a_{n+2} + (n+2) a_n$$

$$\Rightarrow a_{n+2} = \frac{-a_n}{n+1} \text{ is the recurrence relation.}$$

$$a_0 = 0, \quad a_1 = 1, \quad a_2 = \frac{-a_0}{1} = 0, \quad a_3 = \frac{-a_1}{2} = -\frac{1}{2}$$

$$\begin{aligned} \text{Solution is } y &= 0 + 1x + 0x^2 - \frac{1}{2}x^3 + \dots \\ &= x - \frac{1}{2}x^3 + \dots \end{aligned}$$

4. (10 points) The following two equations have a singular point at $x_0 = 0$. One of them has a regular singular point, while the other has an irregular singular point.

$$\text{Equation (A) } x^2 y'' + 4xy' + 2y = 0$$

$$\text{Equation (B) } x^2 y'' + y' + 2y = 0$$

- (a) (4 points) Which equation has a regular singularity, and which has an irregular singularity? Circle your answer below.

(A) is regular, and (B) is irregular. OR (B) is regular, and (A) is irregular.

- (b) (6 points) Justify your answer to the previous part using limits.

$$(A) \text{ is } y'' + \frac{4}{x} y' + \frac{2}{x^2} y = 0$$

$\overset{p(x)}{\underset{||}{x}}$

$\overset{q(x)}{\underset{||}{x^2}}$

$$\lim_{x \rightarrow 0} x p(x) = \lim_{x \rightarrow 0} 4 = 4 \text{ exists and is finite}$$

$$\lim_{x \rightarrow 0} x^2 q(x) = \lim_{x \rightarrow 0} 2 = 2 \text{ exists and is finite}$$


so (A) is regular

$$(B) \text{ is } y'' + \frac{1}{x^2} y' + \frac{2}{x^2} y = 0$$

$\overset{p(x)}{\underset{||}{x^2}}$

$\overset{q(x)}{\underset{||}{x^2}}$

$$\lim_{x \rightarrow 0} x p(x) = \lim_{x \rightarrow 0} \frac{1}{x} \text{ Does not exist}$$



so (B) is irregular.

LAPLACE TRANSFORM

5. (5 points) Write the definition of the Laplace transform of a function $f(t)$.

$$\mathcal{L}\{f(t)\} = \int_0^{\infty} e^{-st} f(t) dt$$

6. (15 points, 5 points per part) Here are some functions of s . Find their inverse Laplace transforms. The answer may have discontinuities or delta functions in it.

(a) $F(s) = \frac{1-2s}{s^2+4s+5}$ $s^2+4s+5 = (s+2)^2+1^2$

$$F(s) = \frac{1-2s}{(s+2)^2+1^2} = \frac{-2(s+2)+5}{(s+2)^2+1^2} = -2 \frac{s+2}{(s+2)^2+1^2} + 5 \frac{1}{(s+2)^2+1^2}$$

$$f(t) = -2 e^{-2t} \cos t + 5 e^{-2t} \sin t$$

(b) $F(s) = \frac{s(e^{-s} - e^{-3s})}{s^2+4} = (e^{-s} - e^{-3s}) G(s)$

where $G(s) = \frac{s}{s^2+4} = \mathcal{L}\{\cos 2t\}$ so $g(t) = \cos 2t$

$$f(t) = u_1(t)g(t-1) - u_3(t)g(t-3)$$

$$= u_1(t) \cos 2(t-1) - u_3(t) \cos 2(t-3)$$

(c) $F(s) = \frac{G(s)}{(s-\pi)^2}$

For this part, $G(s) = \mathcal{L}\{g(t)\}$, and your answer will be in terms of $g(t)$.

$$F(s) = G(s)H(s) \text{ where } H(s) = \frac{1}{(s-\pi)^2}, \text{ so } h(t) = t e^{\pi t}$$

$$f(t) = \int_0^t h(t-\tau)g(\tau) d\tau = \int_0^t (t-\tau) e^{\pi(t-\tau)} g(\tau) d\tau$$

7. (18 points, 6 points per part) A mass on a spring with mass $m = 5$, damping $\gamma = 3$, spring constant $k = 3$, and subject to an external force $g(t)$ obeys the differential equation

$$5y'' + 3y' + 3y = g(t)$$

We impose the initial conditions $y(0) = 2$, $y'(0) = -1$. In each of the scenarios below, solve for $Y(s)$, the Laplace transform of $y(t)$. Do not take the inverse transform of your answer.

- (a) First scenario: No forcing, $g(t) = 0$.

$$\mathcal{L}\{y'\} = sY(s) - y(0) = sY(s) - 2$$

$$\mathcal{L}\{y''\} = s^2Y(s) - sy(0) - y'(0) = s^2Y(s) - 2s + 1$$

$$\text{so } 5(s^2Y(s) - 2s + 1) + 3(sY(s) - 2) + 3Y(s) = 0$$

$$(5s^2 + 3s + 3)Y(s) - 10s - 1 = 0$$

$$Y(s) = \frac{10s + 1}{5s^2 + 3s + 3}$$

- (b) Second scenario: The forcing function $g(t)$ consists of three delta-function impulses at times $t = \pi$, $t = 2\pi$, and $t = 3\pi$:

$$g(t) = \delta(t - \pi) + \delta(t - 2\pi) + \delta(t - 3\pi)$$

$$\mathcal{L}\{g(t)\} = e^{-\pi s} + e^{-2\pi s} + e^{-3\pi s}$$

$$(5s^2 + 3s + 3)Y(s) - 10s - 1 = e^{-\pi s} + e^{-2\pi s} + e^{-3\pi s}$$

$$Y(s) = \frac{10s + 1}{5s^2 + 3s + 3} + \frac{(e^{-\pi s} + e^{-2\pi s} + e^{-3\pi s})}{5s^2 + 3s + 3}$$

(c) Third scenario: The forcing function $g(t)$ is zero up until time $t = \pi$, at which time it switches on to $\sin(t - \pi)$, or in symbols:

$$g(t) = \begin{cases} 0 & \text{if } t < \pi \\ \sin(t - \pi) & \text{if } t \geq \pi \end{cases}$$

$$g(t) = u_{\pi}(t) \sin(t - \pi)$$

$$\mathcal{L}\{g(t)\} = e^{-\pi s} \mathcal{L}\{\sin t\} = e^{-\pi s} \frac{1}{s^2 + 1}$$

$$(5s^2 + 3s + 3)Y(s) - 10s - 1 = e^{-\pi s} \frac{1}{s^2 + 1}$$

$$Y(s) = \frac{10s + 1}{5s^2 + 3s + 3} + \frac{e^{-\pi s}}{(5s^2 + 3s + 3)(s^2 + 1)}$$

FIRST ORDER SYSTEMS

8. (15 points) Find the eigenvalues and eigenvectors of the matrix A , and find the general solution of the system $\mathbf{x}' = A\mathbf{x}$

$$A = \begin{pmatrix} 2 & 0 \\ 1 & -1 \end{pmatrix}$$

$$\det(A - rI) = \det \begin{pmatrix} 2-r & 0 \\ 1 & -1-r \end{pmatrix} = (2-r)(-1-r) - 0 \cdot 1 \\ = (2-r)(-1-r)$$

roots $r_1 = 2$, $r_2 = -1$ are the eigenvalues.

Eigenvector for $r_1 = 2$: $\begin{pmatrix} 0 & 0 \\ 1 & -3 \end{pmatrix} \begin{pmatrix} ? \\ ? \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$

can take $\vec{v}_1 = \begin{pmatrix} 3 \\ 1 \end{pmatrix}$, all eigenvectors are $c_1 \vec{v}_1$ for constant c_1

Eigenvector for $r_2 = -1$: $\begin{pmatrix} 3 & 0 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} ? \\ ? \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$

can take $\vec{v}_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$, all eigenvectors are $c_2 \vec{v}_2$ for const c_2

General solution of $\vec{x}' = A\vec{x}$:

$$\vec{x}(t) = c_1 e^{2t} \begin{pmatrix} 3 \\ 1 \end{pmatrix} + c_2 e^{-t} \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$