

NAME: Solutions

NetID:

MATH 285 E1/F1 Exam 1 (B)

September 19, 2014

Instructor: Pascaleff

Problem	Possible	Actual
1	20	
2	20	
3	20	
4	20	
5	20	
Total	100	


INSTRUCTIONS:

- Do all work on these sheets.
- Show all work.


1. (20 points) Consider the differential equation

$$\frac{dy}{dx} = -xy$$


Which of the following graphs could be a solution curve of this equation? Circle all that apply.




No, slope should be negative in Quadrant I



No slope is positive in Quadrant I whereas $-xy < 0$.



No, since slope at $y=0$ should be zero.



Yes. In fact general solution is $y = Ce^{-\frac{1}{2}x^2}$

$$\left[\begin{array}{l} \frac{dy}{dx} + xy = 0 \quad p(x) = e^{\int x dx} = e^{\frac{1}{2}x^2} \\ \frac{d}{dx} (e^{\frac{1}{2}x^2} y) = 0 \rightarrow e^{\frac{1}{2}x^2} y = C. \end{array} \right]$$

2. (20 points) An object moves along a one-dimensional axis. Its motion is described by a function $x(t)$. It is subjected to an acceleration given by

$$a(t) = 2 + 2\pi \sin(\pi t).$$

Suppose that at $t = 0$, the velocity is zero: $v(0) = 0$. What is the net change in position between $t = 0$ and $t = 1$? That is, what is $x(1) - x(0)$?

$$\frac{dv}{dt} = a(t) = 2 + 2\pi \sin(\pi t)$$

$$v = \int a(t) dt = \int (2 + 2\pi \sin \pi t) dt = 2t - 2\cos(\pi t) + C$$

Find C :

$$0 = v(0) = 0 - 2\cos(\pi \cdot 0) + C = 0 - 2 + C$$

$$2 = C$$

$$\text{So } v(t) = 2t - 2\cos(\pi t) + 2$$

Integrate $v(t)$ from 0 to 1:

$$\begin{aligned} x(1) - x(0) &= \int_0^1 (2t - 2\cos(\pi t) + 2) dt \\ &= \left[t^2 - 2 \frac{\sin(\pi t)}{\pi} + 2t \right]_0^1 \end{aligned}$$

$$= 1 - 2 \frac{\sin(\pi)}{\pi} + 2 - 0 + 2 \frac{\sin(0)}{\pi} - 0$$

$$= 1 + 2 = \boxed{3}$$

3. (20 points) Find the general solution, valid for $x > 0$, of

$$\frac{dy}{dx} = \frac{2y + x^3}{x}$$

Hint: Linear equation, integrating factor.

Write in standard linear form

$$\frac{dy}{dx} = \frac{2y}{x} + x^2 \rightarrow \frac{dy}{dx} - \frac{2}{x}y = x^2$$

$$P(x) = -\frac{2}{x} \quad Q(x) = x^2$$

Integrating factor:

$$\rho(x) = e^{\int P(x) dx} = e^{\int -\frac{2}{x} dx} = e^{-2 \ln(x)}$$

$$= (e^{\ln(x)})^{-2} = x^{-2}$$

Multiply: $x^{-2} \frac{dy}{dx} - 2x^{-1}y = 1$

Product rule $\frac{d}{dx}(x^{-2}y) = 1$

Integrate: $x^{-2}y = x + C$

$$y = x^3 + Cx^2$$

4. (20 points) Consider the equation

$$\frac{dy}{dx} - 2y = xy^2$$

Use the substitution $u = y^{-1}$ to transform this equation into a linear equation for u . Do not solve the resulting equation; the purpose of this problem is merely to transform the original equation for y into one for u .

$$u = y^{-1} \rightarrow y = u^{-1} \rightarrow \frac{dy}{dx} = -u^{-2} \frac{du}{dx}$$

$$\frac{dy}{dx} - 2y = xy^2$$

$$-u^{-2} \frac{du}{dx} - 2u^{-1} = xu^{-2}$$

mult. by $-u^2$

$$\boxed{\frac{du}{dx} + 2u = -x}$$

5. (20 points) A metal ball has been heated to 2000°C . It is placed into a bath of ice water at 0°C . After 10 seconds, it has cooled to a temperature of $(2000e^{-10})^{\circ}\text{C}$ (approximately 0.091°C).

Suppose now that the metal ball is heated again to 2000°C , but instead it is placed into boiling water at 100°C . How long will it take to reach a temperature of 200°C ?

In both situations, the cooling process is governed by Newton's law of cooling:

$$\frac{dT}{dt} = -k(T - A)$$

where A is the temperature of the water, and k is a constant.

First, solve N.L.C. using separation of variables

$$\int \frac{dT}{T-A} = \int -k dt \rightarrow \ln|T-A| = -kt + C$$

$$\rightarrow T-A = \pm e^C e^{-kt} = D e^{-kt}$$

$$\text{so } T = A + D e^{-kt}$$

First situation: $A = 0$ $T(0) = 2000$
 $T(10) = 2000 e^{-10}$

So $2000 = T(0) = 0 + D e^{-k \cdot 0} = D$

Thus $D = 2000$

$$2000 e^{-10} = T(10) = 0 + 2000 e^{-k \cdot 10}$$

$$e^{-10} = e^{-k \cdot 10}$$

$$-10 = -k \cdot 10$$

$$1 = k$$

Thus $k = 1$.

Solution continues



This page is for work that doesn't fit on the other pages. Please indicate the problem that the work goes with.

Second situation: $A=100$ $T(0)=2000$

$$2000 = T(0) = 100 + D e^{-1 \cdot 0}$$

$$2000 = 100 + D$$

Thus $D=1900$.

$$T(t) = 100 + 1900 e^{-t}$$

When is $T(t)=200$?

$$200 = 100 + 1900 e^{-t}$$

$$100 = 1900 e^{-t}$$

$$\frac{1}{19} = e^{-t}$$

$$\ln\left(\frac{1}{19}\right) = -t$$

$$t = -\ln\left(\frac{1}{19}\right)$$

this can be further simplified to $\ln(19)$, but that is not necessary.