

NAME:

NetID:

---

MATH 285 E1/F1 Exam 1 (B)

September 19, 2014

Instructor: Pascaleff

Problem	Possible	Actual
1	20	
2	20	
3	20	
4	20	
5	20	
Total	100	





**INSTRUCTIONS:**

- Do all work on these sheets.
- Show all work.

1. (20 points) Consider the differential equation

$$\frac{dy}{dx} = -xy$$

Which of the following graphs could be a solution curve of this equation? Circle all that apply.



2. (20 points) An object moves along a one-dimensional axis. Its motion is described by a function  $x(t)$ . It is subjected to an acceleration given by

$$a(t) = 2 + 2\pi \sin(\pi t).$$

Suppose that at  $t = 0$ , the velocity is zero:  $v(0) = 0$ . What is the net change in position between  $t = 0$  and  $t = 1$ ? That is, what is  $x(1) - x(0)$ ?

3. (20 points) Find the general solution, valid for  $x > 0$ , of

$$\frac{dy}{dx} = \frac{2y + x^3}{x}$$

*Hint: Linear equation, integrating factor.*

4. (20 points) Consider the equation

$$\frac{dy}{dx} - 2y = xy^2$$

Use the substitution  $u = y^{-1}$  to transform this equation into a linear equation for  $u$ . Do not solve the resulting equation; the purpose of this problem is merely to transform the original equation for  $y$  into one for  $u$ .

5. (20 points) A metal ball has been heated to  $2000^\circ C$ . It is placed into a bath of ice water at  $0^\circ C$ . After 10 seconds, it has cooled to a temperature of  $(2000e^{-10})^\circ C$  (approximately  $0.091^\circ C$ ).

Suppose now that the metal ball is heated again to  $2000^\circ C$ , but instead it is placed into boiling water at  $100^\circ C$ . How long will it take to reach a temperature of  $200^\circ C$ ?

In both situation, the cooling process is governed by Newton's law of cooling:

$$\frac{dT}{dt} = -k(T - A)$$

where  $A$  is the temperature of the water, and  $k$  is a constant.

This page is for work that doesn't fit on the other pages. Please indicate the problem that the work goes with.